

A Construction of Approximately Finite-Dimensional
non-ITPFI Factors

E. J. Woods (Queen's Univ. & RIMS, Kyoto Univ.)

The following is a report on some joint work with
A. Connes which was carried out in Paris in January, 1976.

A von Neumann algebra is said to be approximately finite-dimensional if it is of the form

$$M = \{ M_1 \subset M_2 \subset \dots \subset M_n \subset \dots \}$$

where each M_n is a finite-dimensional matrix algebra. A factor is said to be ITPFI if it is of the form

$$M = \bigotimes_{n=1}^{\infty} (M_n, \omega_n)$$

where each M_n is a type I factor (and ω_n is a state on M_n).

The existence of factors which are approximately finite-dimensional but not ITPFI is an interesting problem. The first construction of such factors was given by Krieger [8]. However in [8] it is only proved that the factors are not "weakly equivalent" to any ITPFI factor. The first proof that these factors are not ITPFI was given by Connes [3]. Alternatively one could now use Krieger's theorem [9] that unitary equivalence implies "weak equivalence" to complete the argument. However Krieger's construction is rather involved, and the arguments of both Krieger [8] and Connes [3] were quite delicate. We give here a new construction for which, in the context of the flow of weights, the argument is rather elementary.

Sec. 1 reviews the relevant aspects of the flow of weights [4], and gives some terminology. Sec. 2 contains the technical lemmas. In Sec. 3 we discuss the examples.

1. Preliminary Material

Let M be a factor, $\text{Aut } M$ the group of all automorphisms of M with the topology of pointwise convergence in the predual, and $\text{Int } M$ the subgroup of inner automorphisms of M . The flow of weights of M is an ergodic action of \mathbb{R}_+^* on some measure space (X_M, μ_M) . The construction of [4] gives not the measure space, but the measure algebra whose elements are unitary equivalence classes $[\phi]$ of integrable weights ϕ of infinite type. The flow is then defined by $F_M(\lambda) [\phi] = [\lambda\phi]$. Let $\alpha \in \text{Aut } M$. The equation $\text{Mod } \alpha [\phi] = [\phi \circ \alpha]$ defines a Borel (and hence continuous) homomorphism from the polish group $\text{Aut } M$ into the polish group of automorphisms of the measure space (X_M, μ_M) . Clearly $\alpha \in \text{Int } M$ implies that $\text{Mod } \alpha = 1$. If M is a factor of type III_0 then the flow of weights $F_{M \otimes M}(\lambda)$ for $M \otimes M$ is given by the action of $F_M(\lambda) \otimes 1$ on the measure algebra of the $F_M(\lambda) \otimes F_M(\lambda^{-1})$ invariant sets on $X_M \times X_M$.

All Borel spaces considered in this paper are standard (i.e. Borel isomorphic to a Borel subset of the unit interval). A transformation S on a measure space (X, μ) is called non-singular if it is invertible and both S and S^{-1} are μ -measurable. Given a non-singular S , the orbit of x under S is the set

$$O_S(x) = \{S^j x : j \in \mathbb{Z}\}.$$

The full group of S is the set $[S]$ of all non-singular transformations T such that for a.e. x , $Tx \in O_S(x)$. A set $W \subset X$ such that $\mu(S^j W \cap S^k W) = 0$ for all $j \neq k$ is called a wandering set for S . S is said to be dissipative if there is a wandering set W such that $X = \bigcup_{j=-\infty}^{\infty} S^j W$.

2. The Technical Lemmas

Let M be a von Neumann algebra, $x, y \in M$. The automorphism σ of $M \otimes M$ defined by the equation $\sigma(x \otimes y) = y \otimes x$ is called the Sakai flip.

Lemma 2.1: Let M be an ITPFI factor, σ the Sakai flip on $M \otimes M$.

Then $\sigma \in \overline{\text{Int}}(M \otimes M)$.

Proof: Let $M = \bigotimes_{n=1}^{\infty} (M_n, \omega_n)$ be given on $\bigotimes_{n=1}^{\infty} (H_n, \Omega_n)$ where

$\omega_n(x) = (x\Omega_n, \Omega_n)$. Then $M \otimes M = \bigotimes_n (M_n \otimes M_n, \omega_n \otimes \omega_n)$ acts on

$K = \bigotimes_n (H_n \otimes H_n, \Omega_n \otimes \Omega_n)$. Let $\psi \in (M \otimes M)_*$, $\varepsilon > 0$.

We can assume that $\bigotimes_n (\Omega_n \otimes \Omega_n)$ is a separating vector for $M \otimes M$ (see lemma 3.15 of [2]). Hence there is a vector $\Psi \in K$ such that

$\psi(x) = (x\Psi, \Psi)$. By lemma 3.1 of [1] there exists $m < \infty$ and $\Psi(m) \in \bigotimes_{n=1}^m (H_n \otimes H_n)$, $\|\Psi(m)\| = 1$, such that $\|\Psi - \Psi_\varepsilon\| < \varepsilon$

where

$$\Psi_\varepsilon = \Psi(m) \otimes \left(\bigotimes_{n=m+1}^{\infty} (\Omega_n \otimes \Omega_n) \right).$$

Let ψ_ε be the state defined by Ψ_ε , and let σ_m be the Sakai flip on $\bigotimes_{n=1}^m (M_n \otimes M_n)$. Then $\sigma\psi_\varepsilon = (\sigma_m \otimes 1)\psi_\varepsilon$. Hence

$$\|(\sigma - \sigma_m \otimes 1)\psi\| < 2\varepsilon.$$

Since σ_m is inner, it follows that $\sigma \in \overline{\text{Int}}(M \otimes M)$. QED.

Lemma 2.2: Let R, S be non-singular transformations on the standard measure space (X, μ) . If S is dissipative and R leaves invariant (modulo μ) all S -invariant measurable sets, then $R \in [S]$.

Proof: We first note that if (E, ν) is a countably separated measure space and $f: E \rightarrow E$ satisfies $f(B) = B$ (modulo ν) for all measurable $B \subseteq E$, then $f(x) = x$ (a.e. ν). Namely let $(B_n)_{n \in \mathbb{N}}$ separate points in E . Then

$$\{x: f(x) \neq x\} \subset \bigcup_n \{B_n \setminus f(B_n)\}$$

which is a set of measure zero.

Now let W be a wandering set for S such that $X = \bigcup_{k=-\infty}^{\infty} S^k W$. Let P_k be the projection of X onto $S^k W$ defined by $P_k x = y$ if $x = S^j y$ for some j such that $y \in S^k W$. Let A be any measurable subset of $S^k W$. Then $\bigcup_{p=-\infty}^{\infty} S^p A$ is S -invariant and it follows that $(P_k R P_k)A = A$ (modulo μ). Now clearly $R \in [S]$ if and only if $P_k R P_k(x) = x$ (a.e.) for all k . **QED.**

The following theorem uses the base and ceiling function construction of a flow. For this purpose it is more convenient to have the flow as an action of \mathbb{R} rather than \mathbb{R}_+^* . Hence we shall use $\mathfrak{F}_M(\lambda) = F_M(e^\lambda)$.

Theorem 2.3: Let M be a factor of type III_0 whose flow of weights can be built under a constant ceiling function with a base transformation T such that $T \times T^{-1}$ is dissipative. Then the Sakai flip $\sigma \notin \overline{\text{Int}(M \otimes M)}$ and hence M is not ITPFI.

Proof: Clearly $\text{Mod } \sigma$ acts on $X_M \times X_M = (B \times I) \times (B \times I)$ by $\sigma(x,s,y,t) = (y,t,x,s)$. Let E be any $T \times T^{-1}$ invariant set in $B \times B$, σ_B the flip on $B \times B$. Then $E \times I \times I$ is an $\mathfrak{F}_M(\lambda) \otimes \mathfrak{F}_M(-\lambda)$ invariant set in $X_M \times X_M$. Now assume $\text{Mod } \sigma = 1$. Then σ_B must preserve E , hence $\sigma_B \in [T \times T^{-1}]$ by the preceding lemma. But this implies that for a.e. $(x,y) \in B \times B$ there exists an integer $n(x,y)$ such that

$$\sigma_B(x,y) = (y,x) = (T^{n(x,y)}x, T^{-n(x,y)}y),$$

i.e. $y \in O_T(x)$. But $O_T(x)$ is countable. QED.

3. The Examples

It remains to demonstrate the existence of approximately finite-dimensional factors of type III_0 satisfying the conditions of Theorem 2.3. For this we first need the existence of ergodic transformations T such that $T \times T^{-1}$ is dissipative. It is a classical result in ergodic theory that such transformations exist [6]. As a specific example, one can use the Markov shift obtained from a two-dimensional random walk. (These transformations preserve an infinite measure.) The existence now follows from the fact that any flow arises as the flow of weights of some approximately finite-dimensional factor [4, 9]. (The proof of this in the general case is not so easy. However for measure preserving flows the argument is not difficult (see for example [7]).)

We remark that $\sigma \in \overline{\text{Int}}(M \otimes M)$ is not a sufficient condition for M to be ITPFI. Namely let M be an approximately finite-dimensional factor whose flow can be built under a constant ceiling function

with a base transformation T which preserves a finite measure. If T is a Bernoulli shift then M is not ITPFI [5]. But then $T T^{-1}$ is ergodic, and it follows easily that $\text{Mod } \sigma = 1$. Hence $\sigma \in \overline{\text{Int}}(M, M)$ [4]. In fact if T is any ergodic transformation preserving a finite measure, it follows from the proof of part (2) of lemma 1 of [7] that $\text{Mod } \sigma = 1$ (see also [10]).

- [1] H. Araki and E. J. Woods, Complete Boolean Algebras of Type I Factors, Publ. Res. Inst. Math. Sci. Ser. A 2 (1966), 157-242.
- [2] H. Araki and E. J. Woods, A classification of factors, Publ. Res. Inst. Math. Sci. Ser. A 4 (1968), 51-130.
- [3] A. Connes, Une classification des facteurs de type III, Ann. Sci. E.N.S. 4eme Serie 6 (1973), 133-252.
- [4] A. Connes and M. Takesaki, Flow of weights on type III factors (to appear)
- [5] A. Connes and E. J. Woods, to appear
- [6] T. E. Harris and H. Robbins, Ergodic theory of Markov chains admitting an infinite invariant measure, Proc. Nat. Acad. Sci. U.S.A. 39 (1953), 860-864.
- [7] T. Hamachi, Y. Oka and M. Osikawa, Flows Associated with Ergodic Non-Singular Transformation Groups, Publ. Res. Inst. Math. Sci. 11 (1975), 31-50
- [8] W. Krieger, On the infinite product construction of nonsingular transformations of a measure space, Invent. Math. 15 (1972), 144-163.
- [9] W. Krieger, On ergodic flows and the isomorphism of factors, Math. Ann. 223 (1976), 19-70.
- [10] T. Hamachi and M. Osikawa, Fundamental Homomorphism of Normalizer Group of Ergodic Transformation, preprint.